## COURS INTENSIF EUROPEEN 19-22 MAI 1992

EUROPEAN INTENSIVE COURSE 19-22 MAY 1992

Prévention des risques d'érosion et de submersion littoraux: la connaissance du risque, les études d'impact en vue des travaux de protection

Prevention of coastal erosion and submersion risks: knowledge of the risk, impact studies with a view to protection works

## Organisé par le Centre Européen sur les Risques Géomorphologiques

Irganised by the European Centre on Geomorphological Hazards

Sous la Direction de Directed by

## Professeur Jean-Claude FLAGEOLLET

Avec le Concours de With the support of

- Conseil de l'Europe: Accord Partiel Ouvert Risques Majeurs
- Ministère de l'Environnement: Délégation aux Risques Majeurs
- Direction Régionale à l'Environnement, Basse- Normandie
   Direction Départementale de l'Equipement du Calvados
- Université de Caen: Laboratoire de Géologie marine

## MARINE EROSION AND LANDSLIDE ACTIVATION AT VILLERVILLE

J. Cl. FLAGEOLLET, Centre Européen sur les Risques Géo- et Morpho-Dynamiques, Strasbourg

either side of it? Above all, however, is phenomena of instability related to lateral extension of the two basins on Can it be affected in the short terme by important issue is the fate of the town. the seaward side, because the most long time been protected by a wall on two basins, Cirque des Graves and delay, or slow down, slippages in the of the slide to prevent further slippages sufficiently increase resistance at the toe either side of Villerville. They do no Fosses du Macre. Villerville has for a protection against marine erosion and from occuring, but they do provide have been laid at the top of the beach on do indeed play the essential role. Rocks shown that water conditions in the slope may result from that? It has been piczometrical level during periods of the slope formations, the higher relation to the water which impregnates role, or only a very subordinate one, in what extent does it play an essential and to facilitate further slippages. But to reduce resistance at the toe of the slide the landslide. In so doing, it helps to sea disperses material from the foot of heavy rain and the movements which In the Villerville-Cricqueboeuf area, the

there any risk of mass movement of the entire slope, including the slope on which the town is built, as is happening in the two basins?

Observation of a natural section at the base of the cliff, between two retaining walls and above the sea wall, has shown that Villerville's sustratum was formed by the filling of a paleovalley with superficial Quaternary formations of the head, sand and loam types. This hypothesis, which was at variance with that previously put forward, namely the existence of a large chalk block underlain by marls, was tested in order to dispel the uncertainties relating to the nature of the town's geological base and evaluate the risk of instability.

To this end, an exploratory borehole with a diameter of 160 mm was drilied by the C.E.T.E. of Rouen in the town centre, on the centre line of the Villerville valley.

This borehole (ST 6), whose head is situated at an altitude of 27 m IGN 69, was drilled to a depth of 20 m, ending in compact blue marls at an altitude of




Fig. 1 Profile of the slope at Villerville

similar. The head is simply much sedimentary beds. However that may be sand structures and mixes the various smal cliff itself with those encountered when Secondly, the difficulty in linking up Flageollet and E. encountered when drilling the borehole that the use of a twist drill destroys drill it is not possible to see whether the with those mapped on the cliff by J.C. matching the various formations surveyed on the present certain problems. Firstly, with a In the schematic geological formations identified are the borehole is due to the facbeen reworked or not the various Helluin (1984) may profile made tormations the valley

approximately 7 m IGN 69 (fig. ۳

The successive formations encountered

0-1 m: fill;

1-2 m :calcareous loess with many small flint splinters

2-9,5m: head with flint splinters, chalk fragments and clay

9,5-19,7 m: reworked Albian sands splinters; with flint fragments and

19,7-20m: compact blue marls Refusal

> 9,5 m in the borehole chalkier between the depth of 7,2 section of the cliff. Ħ.

to problems of instability on either side of the town, owing to lateral extension (Fig. 2) of the area in motion Cirque des Graves) may eventually lead stability and provides reassurances as to subject to instability, are felicitiously glauconitic sand resting on clays and in the two basins (Fosses du Macre and its future. However, further disturbances which both explains Villerville's current replaced by a permeable sandy formation marls as in the case of the two basins blocks accompanied by their cushion of Villerville itself, the hypothetical chalk Quaternary confirms the hypothesis of the filling of and the "Le Bouloir" villa. This fully scarp in the vicinity of the blockhouse Graves: they are visible on the vertical slope (in terms of gradient and direction) nowever, on the side of the Cirque des deposits. Chalk blocks are to be found, of the surfaces of Norman coastal head construction work, closely matches the is built, no doubt partly a result of incline of the slope on which the town the centre line of the valley. The regular No chalk block, was driller through paleovalley deposits. with Hence, superficial




Fig. 2 Zoning of risks of landslides in the Villerville and Cricqueboeuf area