## COURS INTENSIF EUROPEEN 19-22 MAI 1992

EUROPEAN INTENSIVE COURS

# Prévention des risques d'érosion et de submersion littoraux: la connaissance du risque, les études d'impact en vue des travaux de protection

Prevention of coastal erosion and submersion risks: knowledge of the risk, impact studies with a view to protection works

### Organisé par le Centre Européen sur les Risques Géomorphologiques

Organised by the European Centre on Geomorphological Hazards

Sous la Direction de Directed by

## Professeur Jean-Claude FLAGEOLLET

Avec le Concours de With the support of

- Conseil de l'Europe: Accord Partiel Ouvert Risques Majeurs Ministère de l'Environnement: Délégation aux Risques Majeurs
- Direction Régionale à l'Environnement, Basse- Normandie
- Direction Départementale de l'Equipement du Calvados
- Université de Caen: Laboratoire de Géologie marine

# QUATERNARY SLOPE FORMATIONS BETWEEN TROUVILLE AND HONFLEUR.

J.C. FLAGEOLLET, Centre Européen sur les Risques Géo- et Morpho-Dynamiques (CERD), Strasbourg

giving rise to a small "debris" cliff collapsed or slipped. At the bottom of several metres thick which have eolian loam. In the more gentle slope and weathering of the chalk and of and lined irregularly with a quaternary over a height of more than 50 metres cornice formed in the Cenomanian chalk movements formed the basis for the morphological risks of their recent water line, these formations are eroded. everywhere by Quaternary formations the Albian sands and clays are concealed Oxfordian and Kimmeridgian marls and down to the shoreline, the higher slope deposit, made of a mixture of zermos map of Trouville. formations and the superficial The random observation of these the slope, 2 to 3 metres above the high (10 to 12 %) which follows on from it fragments resulting from frost-fracturing the upper third of the slope is a rounded Between Trouville and Honfleur (Fig.1)

## THE QUATERNARY FORMATIONS OF THE SLOPE

## Nature of the formations:

They are, firstly, **chalk blocks** which have collapsed or slipped from the cornice and are generally more voluminous in the intermediate section of the slope (over 100 metres long) than

unconformable Quaternary formations of Kimmeridgian marls dipping towards water's edge, one can see small blocks often weathered and taking on the vertical as a result of pressure from the clay. The chalk always rests on several appearance of a very plastic greenish cracked, sometimes decarbonated, very gritty chalk, always glauconitic, (particularly blocks of chalk). the top of the slope and standing almost their "cushion" of clayey sands. At the the blocks having slipped along with metres of glauconitic clays and sands, featuring large chert beds, jointed, Cenomanian. It is a yellowish sandy or They consist of chalk belonging to the close to the shore (a few dozen metres) "glauconiticy chalk" level of the lower

Secondly, they are **eolian deposits** (loess) and periglacial formations of the debris type. The loess deposits come from the Seine estuary, which was very open at the time of the last Quaternary glaciations. They are sandy, with chalk granules, very calcareous (25 to 30 % CaCo3), with traces of occidation or marmorosis; having been reworked by run-off, they have a slightly laminated structure. The **periglacial debris** inserted between these loess deposits consists of frost-fractures chips of chert and flint and chalk nodules enbedded in more or less sandy (chalk)




Fig. 1 Geomorphology of the Pays d'Auge cliffs

#### chronostratigraphy Tentative formations Development of quaternary

most comprehensive observations can be carried out on the coastal cliff just very deep, they are enlightening. The below Villerville. provided by the recent landslide are not Along the slope, although the sections

cliff reaches a height of + 18 metres known as "Aigue-Marine", where the Section I lies close to the property NGF. Oriented N 50°, this section cuts

## QUATERNARY SLOPE FORMATIONS BETWEEN TROUVILLE AND HONFLEUR

a thin layer covering the chalk blocks and loess deposits. Their presence under a chalk block to a stratified formation of we have been able to see in the bench highly glauconitic clay masses which the chalk blocks is unlikely, despite the observed at the surface, where they form These soliflucted formations are matrix resulting from solifluction types and sizes embedded in a clay formations there are materials of varying addition to these clearly defined they may be 2 to 3 metres thick. In where they have been trapped and where mainly found at the front of the blocks, and tilted forwards. Having accumulated the "head" type with rough cherts graded chalky debris pressed against the wall of on the slope and the changes in its varies greatly according to its location foreshore at the time of the January intumescences which appeared on the from the foot of the cornice, they are form: one goes very quickly from 1982 landslide. panels and the

 $\Box$ 

gravel and non-calcareous

ioams;

comprises three lithostratographic units

Section I starts at + 5 m NGF. It

(fig. 2.I) from bottom to top:

and a height of over 10 metres.

over a length of more than 50 metres valley, which are visible at this point

across the formations of the Villerville


Units II and III represent the same cycle, 日: Weichselian. which may complex rough head likewise, carbonated be placed ij

On the one hand, the head is not chalky

of the glacial period. The traces of On the slope, small blocks of chalk chalky gravel overlain by very thick older cycle, perhaps Saalian. Further up Racine, the underlying loess having equivalent of the brown head of Portis old Weichselian head, perhaps the is not composed of eolian loam enough to be compared with Saalian periods of warming. The lower pedogenesis might be due to interstadial possible been deposited at the beginning period. A more likely theory is that it very start of the upper pleniglacial deposits, it is difficult to place it at the calcareous loess. the valley, the head becomes a very decarbonated unit could be related to an head. On the other, given that its matrix

may be drawn from this: the chalk and loess. They slope gently towards the are incorporated into the head, but the bottom of the valley. Two conclusions large chalk blocks are covered by head

local chalky deposits (sands, silts, undeniable, despite very substantial The eolian origin of the loams is



blocks slid down at a later stage and produced the debris which formed the olocks had largely been completed wher mainly with loess deposits; the nead. It is possible, however, that small rost-fracturing of the chalky cornice he last loess was deposited and when downward movement of the main chalk absence of chalk blocks along the centre valley accounts for the almost complete was smoothed down; the gap in the cornice at the top of the Villerville period; the initial profile of the slope high humidity at the start of the cold disappearance of forest cover and the of the slope and breaking up in the blocks thus sliding down to the bottom underlying glauconitic clays, the chalk ine of process; slippage was facilitated by the blocks set in motion by flowage of the logged Albian sands; this valley, which is filled break-up of

then through increasingly thick head and to West, the fault cuts through chalk: have accumulated. Running from East chalk block tilting towards the West, at corresponding to the back of a huge of the land by three to four metres over the front of which periglacial formations the edge of a shelf-like structure landslide of 1982. The fault occurred at more than 300 metres during the the RN 513) corresponds to the northern Section II (Chemin des Graves, near wall of a fissure which lowered the level

can lead to subsidence and slippage.

during which they slid down on the clay blocks were formed at the beginnin, the cornice due to tapping of water have been as follows : destabilisation of The sequence of events migh and C.8 are probably related to the granules). The two loess deposits C.

#### here. beginning of the Weichselian. Hence, an examination of the cliff solifluction in the early Weichselian and formations and of the kesselt level, which is greatly expanded other two, might be the equivalent of the upper head (C.4) would represent the the loam deposited by periglacial (C.6), which is very different from the the Seine valley. The light brown loam recente (pleniglacial) loess deposits of

very probably moved down at the very Graves suggests that the chalk blocks Chemin des

#### AND LANDSLIDES SLOPE FORMATIONS

### to slidding Susceptibility of formations

were incorporated into the formations.

which, in addtion, often rest on green movement on the slope. The same susceptible to flowage. Heads with a Albian sands. Impregnation with water whose base contains glauconite and cannot be said of the large chalk blocks responsible for a resumption of asserted that these formations are not propelled into motion again. It can be or flints bonded together by their sandy matrix and frost-fractured cherts sandy, are fairly permeable and not very irregular shapes are unlinkely to be The loess deposits, which are very often




Fig. 2 - 2 Section along a secondary scarp at the "chemin des Graves"

### Resumption of movement Holocene Epoch

exactly in how many stages or over extent was underestimated. It is in these slope tend to be located on either side of surveys show that movements of the chaotic appearance. Morphological two areas that one can see blocks of phenomenon, but unfortunately, its what period the phenomenon took place. the Cirque des Graves. One cannot say Villerville, in the Fosses du Macre and due to debris and loess deposits is A slope with a relatively graded profile the slope, allowance was made for this When the villas were built at the top of followed in space and in time by one of

ij

several7different chalk blocks or do not

cut through

chalk tilted and forming a counterslope

clays of Villerville are tilted nearly 45° to be affected by these reverse slopes. with an upslope dip. The cornstone easily removes material, the marls and retreating because the sea attacks it and metres above the beach and rapidly blocks. In the low cliff standing a few fully coincide with the limits of those The resumption of movement has norizontal on the foreshore, also seem limestones of Hennequeville, which are

re-opened several times as a result of proceeded in stages in the Holocene vegetation cover. this have been conceald have appeared successively or have been Epoch, and the small fault scarps which

### J.C. FLAGEOLLET

### REFERENCES

Caen, 125 p. 45 pl., 3 cartes. Esquisse géologique et morphologique. BIGOT A., 1942: La Basse-Normandie.

glissement de terrain, 22 p. Rouen, 1973. Zone côtière entre CENTRE D'ETUDES TECHNIQUES Trouville et Honfleur. Etude de DE L'EQUIPEMENT. Lab. Rég. de

sur les formations de versants DU CNRS CAEN, 1979. Table ronde CENTRE DE GEOMORPHOLOGIE hétérogènes. Bull. n° 24, 200 p.

ROUEN, 1983. Le Quaternaire de DÉPARTEMENT DE GÉOLOGIE DE DU CNRS CENTRE DE GÉOMORPHOLOGIE Normandie, 160 p. DE CAEN ET

Hermann, 241 p., 22 fig., 5 cartes, 7 pl., 7 tabl. Géologie régionale de la France. Paris. DANGEARD L., 1951: La Normandie.

dentale entre la Seine et le Golfe 624p., 136 fig., 31 dépl., 32 pl.h.t. Normand-Breton. Bordeaux. Bière, ELHAI H., 1963: La Normandie occi-

stratigraphique et paléontologique supérieur de la Baie de la Seine : Etude GUYADER J., 1968: Le Jurassique

Thèse, Université. Paris, 268 p.,

d'Auge. Cahier du Département de 3 - 4, Novembre 1971, p. 23-34. Géographie de l'Université de Caen, n° travaux d'aménagement - Le Pays Les terrains du Pays d'Auge et les HELLUIN E. et HELLUIN M., 1971:

geotechnics of coastal cliff stabilisation. London, p. 153-159 HUTCHINSON J. et al., 1982: The

crétacée sur la bordure orientale du Cénomanien de Normandie et du Maine Massif Armoricain; Aptien, Albien, JUIGNET P., 1974: La transgression 806 p., tome 2, 174 fig. Le stratotype du Cénomanien, tome 1.

Normandie. CNRS, Caen, 836 p. Nord-Ouest et plus particulièrement en périglaciaire pléistocène en Europe du LAUTRIDOU J.P., 1984 : Le cycle

contribution à l'étude morphologique de LAUTRIDOU J.P., 1968: Les Sciences Humaines de l'Université de la région. Faculté des Lettres et Romain (Pays de Caux). Leur formations superficielles de Saint-

Carte Zermos de Trouville. Pont MAHIEU J.L. et GUILLOPE P., 1976: l'Evêque. 1/25.000 BRGM, Paris